



Brain aneurysm facts¹



Brain aneurysms are more common than you may think.

35-60
years old

Brain aneurysms are most prevalent in people ages 35 to 60, but can occur in children as well.

1 in 50
people (6.5 million)

in the United States have an unruptured brain aneurysm.

3:2 ratio

Women are more likely than men to have a brain aneurysm.

2x

African Americans are about twice as likely to have a brain aneurysm rupture compared to whites.

Untreated brain aneurysms may have risk of rupture, resulting in what's called hemorrhagic stroke.



Ruptured brain aneurysms account for 3-5% of all new strokes.

500
thousand

Deaths worldwide each year caused by brain aneurysms, and half the victims are younger than 50.



In the United States, a brain aneurysm ruptures every 18 minutes.

~30
thousand

People in the United States suffer a brain aneurysm rupture each year – 8-10 people per 100,000.



1/8 inch

to nearly an inch.
Most aneurysms are small.

20-50%

of all aneurysms rupture.

Treatment Methods Include

(but not limited to):²



Clipping



Coiling



Flow diversion

1. <https://www.bafound.org/about-brain-aneurysms/brain-aneurysm-basics/brain-aneurysm-statistics-and-facts/> (accessed August 2022)

2. Zhao J, Lin H, Summers R, et al. Current treatment strategies for intracranial aneurysms: an overview. *Angiology*. 2018;69(1):17-30

Medtronic

9775 Toledo Way
Irvine, CA 92618
USA
Tel 1-800-716-6700
Fax 763-526-7888

[medtronic.com](https://www.medtronic.com)

UC202106094aEN © 2022 Medtronic. Medtronic, Medtronic logo, and Engineering the extraordinary are trademarks of Medtronic. All other brands are trademarks of a Medtronic company.