



BIS™ Advance monitor for TIVA

# Perfecting your art. Advancing patient care.

Make a difference in post-TIVA outcomes



# Personalize TIVA dosing. Improve outcomes.<sup>1-4</sup>



There are several potential advantages to total intravenous anesthesia (TIVA). But using only routine monitoring during TIVA may increase the risk of unintended awareness or excessively deep anesthesia.<sup>5</sup>

In order to personalize your propofol dosing based on the individual needs of each patient, you need a direct measurement of the propofol effect on your patient's brain. BIS™ Advance monitoring empowers you by accurately reflecting the anesthetic effect on your patient's brain, allowing you to personalize dosing to:

- Improve emergence and recovery times<sup>1-4</sup>
- Reduce postoperative delirium up to 29%<sup>1-5</sup>



# Easy to use. Easy to configure.

The completely redesigned BIS™ Advance monitor gives you:



- A large, high-resolution, touchscreen monitor that's simple to read
- Configurable data and settings so you can see just the information you want
- Color-coordinated data to quickly review readings
- The ability to track total suppression time detected during the procedure
- Built-in troubleshooting guides with information on clinical parameters and data significance

# Proven algorithm. Trusted insight.

With its validated algorithm, BIS™ technology provides meaningful insight so you can personalize dosing throughout a procedure. Studies show that using less anesthetic agent improves outcomes when using TIVA anesthetic approach, when following protocols supporting adoption of Enhanced Recovery after Surgery guidelines, and in elderly patient populations at risk for postoperative neurocognitive disorders.<sup>2,12</sup> In addition, brain monitoring is recommended in multiple society guidelines.<sup>8,15,16</sup>

# Easy output. Easy to move.

The BIS™ Advance monitor is engineered to help make your workflow more efficient while giving you the insight you need.

- Save time with data output protocols that enable connectivity to electronic medical records (EMRs)
- Maintain continuous monitoring when moving between care settings



To request a full clinical demo of the BIS™ Advance monitor, contact your Medtronic representative.



The BIS™ monitoring system should not be used as the sole basis for diagnosis or therapy and is intended only as an adjunct in patient assessment. Reliance on the BIS™ monitoring system alone for intraoperative anesthetic management is not recommended.

- Whitlock E, Vannucci A, Avidan M. Postoperative Delirium. *Minerva Anestesiol.* 2011 April; 77(4): 448-456.
- Sanders R, Pandharipande P, Davidson A, et al. Anticipating and managing postoperative delirium and cognitive decline in adults. *BMJ.* 2011; 343:d4331.
- Abelha FJ, Luis C, Veiga D, et al. Outcome and quality of life in patients with postoperative delirium during an ICU stay following major surgery. *Crit Care.* 2013;17(5):R257.
- Rudolph J, Marcantonio E. Postoperative Delirium: Acute change with long-term implications. *Anesth Anal.* 2011; 112(5),1202-1211.
- Punjasawadwong Y, Boonjeungmonkol N, Phongchiewboon A. Bispectral index for improving anaesthetic delivery and postoperative recovery. *Cochrane Database Syst Rev.* 2014;(6):CD003843.
- Zywiell MG, Hurley RT, Perruccio AV, et al. Health economic implications of perioperative delirium in older patients after surgery for a fragility hip fracture. *J Bone Joint Surg Am.* 2015; 97(10):829-836.
- Gleason LJ, Schmitt EM, Kosar CM, et al. Effect of delirium and other major complications on outcomes after elective surgery in older adults. *JAMA Surg.* 2015 Sep 9:1-7.
- Inouye SK, Sharon K, et al. Postoperative Delirium in Older Adults: Best Practice Statement from the American Geriatrics Society. Intraoperative Measures to Prevent Delirium. *J Am Coll Surg.* 2014; 220(2):136-148.e1.
- Veiga D, Luis C, Parente D, et al. Postoperative delirium in intensive care patients: risk factors and outcome. *Rev Bras Anesthesiol.* 2012;62(4):469-483.
- Koster S, Hensens AG, Schuurmans MJ, et al. Consequences of delirium after cardiac operations. *Ann Thorac Surg.* 2012; 93(3):705-711.
- Sieber FE, Zakriya K, Gottschalk A, et al. Sedation depth during spinal anesthesia and the development of postoperative delirium in elderly patients undergoing hip fracture repair. *Mayo Clin Proc.* 2010; 85(1), 18-26.
- Chan M, Cheng B, Lee T, et al. BIS-guided anesthesia decreases postoperative delirium and cognitive decline. *Journal of Neurosurgical Anesthesiology.* 2013; 25(1), 33-42.
- Radtke FM, Franck M, Lendner J, et al. Monitoring depth of anaesthesia in a randomized trial decreases the rate of postoperative delirium but not postoperative cognitive dysfunction. *Br J Anaesth.* 2013; 110: i98-i105.
- Zhang C, Xu L, Ma Y-Q, et al. Bispectral index monitoring prevent awareness during total intravenous anesthesia:a prospective, randomized, double-blinded, multi-center controlled trial. *Chin Med J (Engl).* 2011;124(22):3664-3669.
- Checketts M, Alladi R, Ferguson K., et al. Recommendations for standards of monitoring during anaesthesia and recovery. 2015: Association of Anaesthetists of Great Britain and Ireland. *Anaesthesia.* 2016;71(1):85-93. doi: 10.1111/anae.13316.
- Nunes R, Fonseca N, Simões C., et al. Brazilian consensus on anesthetic depth monitoring. *Braz J Anesthesiol.* 2015;65(6):427-436. doi: 10.1016/j.bjane.2015.10.001.

Important: Please refer to the package insert for complete instructions, contraindications, warnings and precautions.

© 2023 Medtronic. Medtronic, Medtronic logo, and Engineering the extraordinary are trademarks of Medtronic.

™™ Third-party brands are trademarks of their respective owners. All other brands are trademarks of a Medtronic company.

EMEA-PM-2200095-bis-advance-monitor-tiva-brochure-en-we-8339237